Scheme for matching historical Land Utilisation Survey (LUS) land-use categories to modern Land Cover Map 2007 (LCM2007) land-cover classes. Classes from the LCM2007 appear in the left-hand columns, with the (LUS) classes they were mapped onto in the third column. There were some cases where the delineation between particular classes was unclear, particularly around the coastline, and so some were either removed or amalgamated (fourth column).

CEH LCM 2007 classes (n = 23)	CEH ID in 25m raster	LUSGB 1933-49 land use classes (n = 8)	Our broad land use categories $(n = 6)$
Urban	22	Urban	Urban
Suburban	23	Suburban	
Broadleaved woodland	1	Woodland	Woodland
Coniferous woodland	2		
Arable and horticulture	3	Arable	Arable
Improved grassland	4	Not mapped	Agriculturally-improved grassland
Neutral grassland	6	Grassland	Grassland
Calcareous grassland	7		
Fen, marsh and swamp	9		
Rough grassland	5	Rough grazing	
Acid grassland	8		
Heather	10		
Heather grassland	11		
Bog	12		
Montane habitats	13		
Inland rock	14		
Saltmarsh	21	Delineation unclear (Removed) (Removed)	(Removed)
Saltwater	15		(Kellioved)
Freshwater	16	Water	Water
Littoral rock	19	Unpredictably assigned to the above (Removed)	(Removed)
Littoral sediment	20		
Supra-littoral rock	17		
Supra-littoral sediment	18		